

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

BLACK BORDERS


- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Problem Image Mailbox.**


Claims

1. An aqueous liquid detergent composition comprising a long-chain fatty acid glycol ester represented by the following formula (1):



wherein, R^1 represents a C_{11-21} alkyl or alkenyl group, A represents a hydrogen atom or COR^2 (in which R^2 represents a C_{11-21} alkyl or alkenyl group), and m stands for a number of from 1 to 3, and a surfactant, wherein the fatty acid
10 constituent of the long-chain fatty acid glycol ester contains 60 wt.% or greater of fatty acids having 18 or more carbon atoms, 40 wt.% or less of fatty acids having 16 carbon atoms, and 5 wt.% or less of fatty acids having less than 16 carbon atoms; and wherein the composition has a pH
15 of from 1 to 5 at 25°C when diluted to 20 times the weight with water.

2. An aqueous liquid detergent composition comprising a long-chain fatty acid glycol ester represented by the following formula (1):



wherein, R^1 represents a C_{11-21} alkyl or alkenyl group, A represents a hydrogen atom or COR^2 (in which R^2 represents a C_{11-21} alkyl or alkenyl group), and m stands for a number of from 1 to 3, and a surfactant, wherein the fatty acid
25 constituent of the long-chain fatty acid glycol ester

contains 65 to 85 wt.% of fatty acids having 18 or more carbon atoms, 15 to 35 wt.% of fatty acids having less than 18 carbon atoms, 35 wt.% or less of fatty acids having 16 carbon atoms, and 5 wt.% or less of fatty acids having less than 16 carbon atoms; and wherein the composition has a pH of from 1 to 5 at 25°C when diluted to 20 times the weight with water.

3. The composition according to Claim 1, wherein the long-chain fatty acid glycol ester of formula (1) is selected from the group consisting of ethylene glycol mono-fatty acid esters, ethylene glycol di-fatty acid esters, triethylene glycol mono-fatty acid esters, triethylene glycol di-fatty acid esters, and mixtures thereof.

4. The composition according to Claim 1 comprising from 0.3% to 10wt% of the composition the long-chain fatty acid glycol ester of formula (1).

5. The composition according to Claim 1 wherein the surfactant is selected from the group consisting of amphoteric surfactants, nonionic surfactants, and mixtures thereof.

6. The composition according to Claim 5 comprising from 10 to 50wt. % of surfactant.

7. The composition according to Claim 1 further comprising from 0.05% to 10 wt.% of an organic acid or inorganic acid.

8. The composition according to Claim 3, wherein the long-chain fatty acid glycol ester of formula (1) is an ethylene glycol di-fatty acid ester.

5 9. The composition according to Claim 1, further comprising a silicone selected from the group consisting of dimethylpolysiloxanes, amino-modified silicones, polyether-modified silicones, methylphenylpolysiloxanes, fatty acid-modified silicones, alcohol-modified silicones, alkoxy-modified silicones, epoxy-modified silicones, fluorine-
10 modified silicones, cyclic silicones, alkyl-modified silicones, and mixtures thereof.

10. The composition according to Claim 9, comprising from 0.01 to 10 wt.% of dimethylpolysiloxanes.